arXiv:1704.08444v1 [stat.ME] 27 Apr 2017

A note on Quantile curves based bivariate reliability concepts

Sreelakshmi N and Sudheesh K Kattumannil

Indian Statistical Institute, Chennai, India

Abstract

Several attempts were made in the literature to generalize univariate reliability concepts to bivariate as well as multivariate set up. Here we extend the univariate quantile based reliability concepts to bivariate case based on quantile curves. We propose quantile curves based bivariate hazard rate and bivariate mean residual life function and study their uniqueness properties to determine the underlying quantile curve. A relationship between them is also derived.

1 Introduction

A probability distribution of a continuous random variable X can be represented either in terms of distribution function $F(x)$ or by the quantile function defined by

$$Q_X(u) = \inf\{x : F(x) \geq u\};\ 0 \leq u \leq 1.$$

Gilchrist (2000) discussed various properties and uses of quantile functions in statistical modeling. Nair et al. (2013) proposed univariate reliability concepts in a quantile frame work and studied the properties in detail.

Many researchers generalised the concepts based on univariate quantile functions to bivariate as well as multivariate set up. Tukey (1977) defined the concept of depth function using the characterizations based on minimizing distances which is followed by the proposal of the concept, zonoid quantile by Koshevoy and Mosler (1997) and median balls by Averous and Meste (1997). Later multivariate quantiles are defined as half

planes by Nolan (1992) and Massé and Theodorescu (1994). A series of papers appeared about the generalization of univariate quantile which include Koltchinskii (1997), Chen and Welsh (2002), Mosler (2002) and Serfling (2002).

Fernández-Ponce and Suarez-Llorens (2002) defined the multivariate quantile as a set of points which accumulate the same probability for a fixed orthant known as quantile curves. Belzunce et al. (2007) used this concept of quantile curves to study the dependence structure of bivariate distribution functions. For this purpose, the authors used the concept of copula and we need some definitions and notations to discuss this idea, which is explained below. Let $\mathbf{X} = (X, Y)$ be a bivariate absolutely continuous random vector and $\mathbf{x} = (x, y)$ be a point in R^2. Let $F(.)$ and $G(.)$ be the marginal distribution functions of X and Y respectively. Also let $F_\varepsilon(\mathbf{x})$ be the joint distribution function of $\mathbf{X}$. Denote the four directions in the two dimensional plane as $\varepsilon = (\varepsilon_1, \varepsilon_2)$ with $\varepsilon_i \in \{-1, 1\}; i = 1, 2$. To simplify the notation, we use $-$ and $+$ to represent -1 and $+1$, respectively. For example,

$$F_{\varepsilon_{--}}(\mathbf{x}) = P[X \leq x, Y \leq y].$$

The p^{th} bivariate quantile curve for the direction ε denoted by $Q_{\mathbf{X}}(p, \varepsilon)$ defined as

$$Q_{\mathbf{X}}(p, \varepsilon) = \left\{(\mathbf{x}) \in R^2 : F_\varepsilon(\mathbf{x}) = p\right\}, \quad 0 \leq p \leq 1.$$

As an example, for the random vector $\mathbf{X}$ having bivariate Pareto distribution with independent components following univariate Pareto distribution with scale parameter k and shape parameter α, it is possible to plot four quantile curves in four directions for each $p \in [0, 1]$. For the direction ε_{++} we can get the quantile curve by plotting $xy = k^2 p^{-1/\alpha}$ for $0 \leq p \leq 1$.

Sklar (1959) introduced the concept of copula which is a bivariate distribution function with uniform marginals on $[0, 1]$. For the joint distribution function F_ε with margins F and G, there exists a copula C such that

$$F_\varepsilon(\mathbf{x}) = C(F(x), G(y)).$$

Suppose $F(x) = u$ and $G(y) = v$, then $x = F^{-1}(u) = Q_X(u)$ and $y = G^{-1}(v) = Q_Y(v)$. Therefore $F_\varepsilon\left(Q_X(u), Q_Y(v)\right)$ depends only on the copula C for each direction ε and is obtained as (Belzunce et al., 2007)

$$F_\varepsilon\left(Q_X(u), Q_Y(v)\right) = \begin{cases} C(u,v); & when\ \varepsilon = \varepsilon_{--} \\ u - C(u,v); & when\ \varepsilon = \varepsilon_{-+} \\ v - C(u,v); & when\ \varepsilon = \varepsilon_{+-} \\ 1 - u - v + C(u,v); & when\ \varepsilon = \varepsilon_{++}, \end{cases}$$

where $0 \leq u \leq 1$ and $0 \leq v \leq 1$.

Sreelakshmi (2017) introduced the bivariate reliability concepts using dependence structure and studied properties as well as characterizations based on the relationship between copula based bivariate hazard rate and bivariate mean residual life. For each direction, the probability $F_\varepsilon(.)$ is written in terms of copula and thus lead to the construction of bivariate reliability concepts. For the bivariate random vector $\mathbf{X} = (X, Y)$, Belzunce et al. (2007) showed that the quantile curves can be expressed by means of quantiles of the conditional distributions of $Y|X \leq x$ and $Y|X \geq x$ given by

$$\begin{aligned} Q_{\mathbf{X}}(p, \varepsilon_{--}) &\to \left\{ \left(Q_X(u), Q_{Y|X \leq Q_X(u)}(\frac{p}{u}) \right) : u > p \right\} \\ Q_{\mathbf{X}}(p, \varepsilon_{+-}) &\to \left\{ \left(Q_X(u), Q_{Y|X \geq Q_X(u)}(\frac{p}{1-u}) \right) : u < 1 - p \right\} \\ Q_{\mathbf{X}}(p, \varepsilon_{-+}) &\to \left\{ \left(Q_X(u), Q_{Y|X \leq Q_X(u)}(1 - \frac{p}{u}) \right) : u > p \right\} \\ Q_{\mathbf{X}}(p, \varepsilon_{++}) &\to \left\{ \left(Q_X(u), Q_{Y|X \geq Q_X(u)}(1 - \frac{p}{1-u}) \right) : u < 1 - p \right\}. \end{aligned} \tag{1}$$

where $0 \leq u, p \leq 1$.

In short, the four level curves can be expressed in terms of dependence structure (using the concept of copula) or by means of quantiles of conditional distributions. It is interesting to note that the second way of expressing the level curve does not require any dependence measure and it requires only the quantiles of conditional distributions.

Motivated by the above fact, in the present work an attempt is made to study the reliability concepts in bivariate set up using quantiles of conditional distributions. The extension of univariate quantile based reliability concepts to the bivariate set up can

be done on the basis of the four directions. Here we consider the bivariate reliability concepts in the direction ε_{--}. The other directions can be studied in similar fashion. Section 2 gives the definitions on quantile curves based bivariate hazard rate and bivariate mean residual life function and their uniqueness properties and relationship are explained in Section 3.

2 Quantile curves based bivariate reliability concepts

To introduce the bivariate reliability concepts based on quantile curves, we need some notations which are explained below. Let $\mathbf{X} = (X, Y)$ be the bivariate random vector with univariate marginal distribution function $F(.)$ and $G(.)$. Suppose that $F(x) = u$ so that $x = F^{-1}(u) = Q_X(u)$ and $G(y) = v$ so that $y = G^{-1}(v) = Q_Y(v)$. Let $Y_X = Y|X \leq x$ and $G_1(y) = P(Y \leq y|X \leq x) = P_X$. Therefore, $Y_X = G_1^{-1}(P_X) = \phi(P_X)$. Also $X_Y = X|Y \leq y$ and $G_2(x) = P(X \leq x|Y \leq y) = P_Y$ which yields $X_Y = G_2^{-1}(P_Y) = \psi(P_Y)$.

Next we propose the definitions for bivariate hazard rate as well as bivariate mean residual life based on quantile curves for the direction ε_{--}. From (1), the quantile curve in the direction ε_{--}, $Q_{\mathbf{X}}(p, \varepsilon_{--})$ is a vector containing u^{th} quantile corresponding to the random variable X denoted as $Q_X(u)$ and $Q_{Y|X \leq Q_X(u)}(p/u)$ which is the quantile function corresponds to the random variable Y_X and a point that accumulates probability $\frac{p}{u}$ to the left tail and $1 - \frac{p}{u}$ to the right tail provided $u > p$. Based on the above vector of two quantile functions, we look into the definitions of quantile curves based bivariate reliability concepts.

Definition 1. *The bivariate hazard rate in the direction ε_{--} in terms of quantile curves can be defined as the vector*

$$\underset{\varepsilon_{--}}{h}(u, P_X) = (h_1(u), h_2(P_X)) \tag{2}$$

with

$$h_1(u) = \frac{1}{(1-u)Q'_X(u)} \quad \textit{and} \quad h_2(P_X) = \frac{1}{(1-P_X)\phi'(P_X)}, \tag{3}$$

where prime denotes the differentiation.

Apart from the hazard rate, it is of much interest in studying the concept of remaining life time of a unit given it has survived up to a particular time point t. To know about the definition and properties of bivariate mean residual life, one can refer to Nair and Nair (1989) and Kulkarni and Rattihalli (1996). Here we give definition of the quantile curves based bivariate mean residual life function.

Definition 2. *The quantile curves based bivariate mean residual life function for the direction ε_{--} as*

$$\underset{\varepsilon_{--}}{m}(u, P_X) = (m_1(u), m_2(P_X)), \tag{4}$$

where

$$m_1(u) = \frac{1}{1-u}\int_u^1 Q_X(z)dz - Q_X(u) \tag{5}$$

and

$$m_2(P_X) = \frac{1}{1-P_X}\int_{P_X}^1 \phi(z)dz - \phi(P_X). \tag{6}$$

Remark 1. *If we interchange X and Y, we obtain another vector for the bivariate hazard rate in the direction ε_{--} and is given by*

$$\underset{\varepsilon_{--}}{H}(v, P_Y) = (H_1(v), H_2(P_Y)) \tag{7}$$

where

$$H_1(v) = \frac{1}{(1-v)Q'_Y(v)} \quad \textit{and} \quad H_2(P_Y) = \frac{1}{(1-P_Y)\psi'(P_Y)}.$$

Also the bivaraite mean residual life is given by

$$\underset{\varepsilon_{--}}{M}(v, P_Y) = (M_1(v), M_2(P_Y)), \tag{8}$$

where

$$M_1(v) = \frac{1}{1-v}\int_v^1 Q_Y(z)dz - Q_Y(v)$$

and

$$M_2(P_Y) = \frac{1}{1-P_Y} \int_{P_Y}^{1} \psi(z)dz - \psi(P_Y).$$

3 Properties and Characterizations

In this section we study the uniqueness property of the concepts derived in previous section. Also we look into the relationship between the bivariate hazard rate and bivariate mean residual life in the quantile curves based approach. Next two theorems provide the uniqueness properties of quantile curves based bivariate hazard rate and bivariate mean residual life. We omit the proof of theorems stated in this section since the derivation is similar to that of univariate quantile based reliabilty concepts given in Nair et al. (2013).

Theorem 1. *For the bivariate random vector* $\boldsymbol{X}$*, quantile curve based bivariate hazard rate defined by (2) determines the underlying quantile curve* $Q_{\boldsymbol{X}}(p, \varepsilon_{--})$ *uniquely. Here* $Q_{\boldsymbol{X}}(p, \varepsilon_{--})$ *is obtained as*

$$Q_{\boldsymbol{X}}(p, \varepsilon_{--}) = \left(Q_X(u), \phi(P_X)\right),$$

where

$$Q_X(u) = \int_0^u \frac{dz}{(1-z)h_1(z)}$$

and

$$\phi(P_X) = \int_0^{P_X} \frac{dz}{(1-z)h_2(z)}.$$

Theorem 2. *For the bivariate random vector* $\boldsymbol{X}$*, the quantile curve can be expressed uniquely in terms of* $\underset{\varepsilon_{--}}{m}(u, P_X)$ *through*

$$Q_{\boldsymbol{X}}(p, \varepsilon_{--}) = \left(Q_X(u), \phi(P_X)\right),$$

where

$$Q_X(u) = \mu_X - m_1(u) + \int_0^u \frac{m_1(z)}{1-z}dz$$

and

$$\phi(P_X) = \mu_{Y_X} - m_2(P_X) + \int_0^{P_X} \frac{m_2(z)}{1-z} dz,$$

and μ_X *and* μ_{Y_X} *are the means corresponding to the random variables* X *and* Y_X*, respectively.*

The quantile curve based bivariate hazard rate obtained after interchanging X and Y given in (8) uniquely determines the underlying quantile curve through

$$Q_Y(v) = \int_0^v \frac{dz}{(1-z)H_1(z)}$$

and

$$\psi(P_Y) = \int_0^{P_Y} \frac{dz}{(1-z)H_2(z)}.$$

Similarly the bivariate mean residual life given in (9) determines the quantile curve through

$$Q_Y(v) = \mu_Y - M_1(v) + \int_0^v \frac{M_1(z)}{1-z} dz$$

and

$$\psi(P_Y) = \mu_{X_Y} - M_2(P_Y) + \int_0^{P_Y} \frac{M_2(z)}{1-z} dz,$$

where μ_Y and μ_{X_Y} are the means of the random variables Y and X_Y respectively.

Theorem 3. *For the bivariate random vector* $\boldsymbol{X}$*, the quantile curves based bivariate hazard rate and bivariate mean residual life function are related through the relationship*

$$(1-u)m_1(u) = \int_u^1 \frac{dz}{h_1(z)}$$

and

$$(1-P_X)m_2(P_X) = \int_{P_X}^1 \frac{dz}{h_2(z)}.$$

Remark 2. *The above proposed quantile curves based bivariate reliability concepts can be also done in reversed time setup. This yields the definitions of bivariate reversed hazard rate and bivariate reversed mean residual life function in a quantile curves based approach.*

References

1. Avérous, J. and Meste, M. (1997), Median balls: an extension of the interquantile intervals to multivariate distributions, *Journal of Multivariate Analysis*, 63, 222-241.

2. Belzunce, F., Castaño, A., Olvera-Cervantes, A. and Suárez-Llorens, A. (2007), Quantile curves and dependence structure for bivariate distributions, *Computational Statistics and Data Analysis*, 51, 5112-5129.

3. Chen, L. A. and Welsh, A. H. (2002), Distribution-function based bivariate quantile, *Journal of multivariate analysis*, 83, 208-231.

4. Fernández-Ponce, J. M. and Suárez-Llorens, A. (2003), A multivariate dispersion ordering based on quantile more widely separated, *Journal of Multivariate Analysis*, 85, 40-53.

5. Gilchrist, W. (2000), *Statistical modeling with quantile functions*, CRC Press, Florida.

6. Koltchinskii, V. I. (1997), M-estimation, convexity and quantile, *Annals of Statistics*, 25, 435-477.

7. Koshevoy, G. and Mosler, K. (1997), Zonoid trimming for multivariate distributions, *Annals of Statistics*, 25, 1998-2017.

8. Kulkarni, H. V. and Rattihalli, R. N. (1996), On Characterization of Bivariate Mean Residual Life Function, *IEEE Transactions on Reliability*, 38, 362-364.

9. Massé, J. C. and Theodorescu, R. (1994), Halfplane trimming for bivariate distributions, *Journal of Multivariate Analysis*, 48, 188-202.

10. Mosler, K. (2002), *Multivariate dispersion, central regions and depth: the lift zonoid approach*, Springer-Verlag, New York.

11. Nair, K. R. M. and Nair, N. U. (1989), Bivariate Mean Residual Life, *IEEE Transactions on Reliability*, 38, 362-364.

12. Nair, N. U., Sankaran, P. G. and Balakrishnan, N. (2013), *Quantile based reliability analysis*, Birkhäuser, Basel.

13. Nolan, D. (1992), Asymptotics for multivariate trimming, *Stochastic processes and their applications*, 42, 157-169.

14. Serfling, R. (2002), Quantile functions for multivariate analysis: approaches and applications, *Statistica Neerlandica*, 56, 214-232.

15. Sklar, A. (1959), Fonctions de reparition a n dimensions et leurs marges, *Publications de l'Institut de Statistique de l'Universite de Paris*, 8, 229-231.

16. Sreelakshmi, N (2017), An introduction to copula based bivariate reliability concepts, *Communication in Statistics-Theory and methods*, accepted manuscript; http://dx.doi.org/10.1080/03610926.2017.1316396.

17. Tukey, J. W. (1977), *Exploratory data analysis*, Addison-Wesley, Princeton.